arXiv:1612.01116v1 [math.FA] 4 Dec 2016

CHARACTERIZATION OF CMO VIA COMPACTNESS OF THE COMMUTATORS OF BILINEAR FRACTIONAL INTEGRAL OPERATORS

DINGHUAI WANG, JIANG ZHOU* AND WENYI CHEN

Abstract. Let I_α be the bilinear fractional integral operator, B_α be a more singular family of bilinear fractional integral operators and $\vec{b}=(b,b)$. Bényi et al. in [1] showed that if $b\in$ CMO, the BMO-closure of $C_c^\infty(\mathbb{R}^n)$, the commutator $[b,B_\alpha]_i(i=1,2)$ is a separately compact operator. In this paper, it is proved that $b\in$ CMO is necessary for $[b,B_\alpha]_i(i=1,2)$ is a compact operator. Also, the authors characterize the compactness of the **iterated** commutator $[\Pi\vec{b},I_\alpha]$ of bilinear fractional integral operator. More precisely, the commutator $[\Pi\vec{b},I_\alpha]$ is a compact operator if and only if $b\in$ CMO.

1. Introduction

A locally integrable function f is said to belong to BMO space if there exists a constant $C>0$ such that for any cube $Q\subset\mathbb{R}^n$,

$$\frac{1}{|Q|}\int_Q |f(x)-f_Q|dx\le C,$$

where $f_Q=\frac{1}{|Q|}\int_Q f(x)dx$ and the minimal constant C is defined by $\|f\|_*$. There are a number of classical results that demonstrate BMO functions are the right collections to do harmonic analysis on the boundedness of commutators. A well known result of Coifman, Rochberg and Weiss [8] states that the commutator

$$[b,T](f)=bT(f)-T(bf)$$

is bounded on some L^p, $1<p<\infty$, if and only if $b\in$ BMO, where T be the classical Calderón-Zygmund operator. In 1978, Uchiyama [21] refined the boundednss results on the commutator to compactness. This is a achieved by requiring the commutator with symbol to be in CMO, which is the closure in BMO of the space of C^∞ functions with compact

2010 Mathematics Subject Classification. Primary 42B20, 47B07; Secondary: 42B25,47G99.

Key words and phrases. Bilinear Fractional Integral Operators; Characterization; Compactness; Iterated Commutator

The research was supported by National Natural Science Foundation of China (Grant No.11661075 and No.11261055).

* Corresponding author, Email: zhoujiangshuxue@126.com.

support. In recent years, the compactness of commutators has been extensively studied already, as Chen, Ding and Wang [5], [6] and Wang [22]. The interest in the compactness of $[b, T]$ in complex analysis is from the connection between the commutators and the Hankel-type operators. In fact, the authors of [13] and [14] have applied commutator theory to give a compactness characterization of Hankel operators on holomorphic Hardy spaces $H^2(D)$, where D is a bounded, strictly pseudoconvex domain in $\mathbb{C}^n$. It is perhaps for this important reason that the compactness of $[b, T]$ attracted ones attention among researchers in PDEs.

In the multilinear setting, the boundedness results for commutators with symbols in BMO started to receive attention only a few years ago, see [15], [17], [18] or [20]. Compactness results in the multilinear setting have just began to be studied. Bényi and Torres [3], Bényi et al. [1] and [2] showed that symbols in CMO again produce compact commutators. Specially, Bényi et al. in [1] showed that if $b \in \mathrm{CMO}$, the commutator $[b, B_\alpha]_i (i = 1, 2)$ is a separately compact operator. More precisely, it is obtained that if $b \in \mathrm{CMO}$ and $g \in L^{p_2}$ is fixed, $[b, B_\alpha]_i(\cdot, g)(i = 1, 2)$ is a compact operator from L^{p_1} to L^q. Unfortunately, it is unknown that if $b \in \mathrm{CMO}$, are the commutators $[b, B_\alpha]_i, i = 1, 2$, jointly compact? We intend to study this question in future work, however, in this paper, we first give the necessary condition for commutators $[b, B_\alpha]_i$ are jointly compact.

Another subject of this paper is to consider the characterization of compactness of the iterated commutator of I_α. In 2015, Chaffee and Torres [4] characterized the compactness of the **linear commutators** of bilinear fractional integral operators acting on product of Lebesgue spaces. In this paper, the characterization of compactness of the **iterated commutators** will be considered.

To state the main result of this paper, we first recall some necessary notions and notation.

It is well known that the fractional integral $\mathcal{I}_\alpha$ of order $\alpha (0 < \alpha < n)$ plays an important role in harmonic analysis, PDE and potential theory (see [19]). Recall that $\mathcal{I}_\alpha$ is defined by

$$\mathcal{I}_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-\alpha}} dy.$$

For the bilinear case, the bilinear fractional integral operator I_α, $0 < \alpha < 2n$, is defined by

$$I_\alpha(f_1, f_2)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f_1(y_1) f_2(y_2)}{\big(|x - y_1| + |x - y_2|\big)^{2n-\alpha}} dy_1 dy_2.$$

In this paper, we will consider the following equivalent operator

$$I_\alpha(f_1,f_2)(x)=\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{f_1(y_1)f_2(y_2)}{\left(|x-y_1|^2+|x-y_2|^2\right)^{n-\alpha/2}}dy_1dy_2.$$

Its iterated commutators with $\vec{b}=(b_1,b_2)$ is given by

$$I_\alpha(f_1,f_2)(x)=\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{(b_1(x)-b_1(y_1))(b_2(x)-b_2(y_2))f_1(y_1)f_2(y_2)}{\left(|x-y_1|^2+|x-y_2|^2\right)^{n-\alpha/2}}dy_1dy_2.$$

We will now examine a more singular family of bilinear fractional integral operators,

$$B_\alpha(f,g)(x)=\int_{\mathbb{R}^n}\frac{f(x-y)g(x+y)}{|y|^{n-\alpha}}dy.$$

This operator was first introduced by Grafakos in [9], and later studied by Grafakos and Kalton [10] and Kenig and Stein [12]. The commutators $[b,B_\alpha]_i$ of B_α with b can be written as

$$\begin{aligned}[b,B_\alpha]_1(f,g)(x) &= bB_\alpha(f,g)-B_\alpha(bf,g)\\ &= \int_{\mathbb{R}^n}\frac{b(x)-b(y)}{|x-y|^{n-\alpha}}f(y)g(2x-y)dy,\end{aligned}$$

the definition of $[b,B_\alpha]_2$ be the similar as $[b,B_\alpha]_1$. In what follows, we need only consider one of these two commutators.

For $1<p\le q<\infty$, recall that the Muckenhoupt class of weights consists of all nonnegative, locally integrable functions ω such that

$$\sup_Q\left(\frac{1}{|Q|}\int_Q\omega^q(x)dx\right)\left(\frac{1}{|Q|}\int_Q\omega(x)^{-p'}dx\right)^{q/p'}<\infty.$$

We also recall the definition of the multiple or vector weights used in the bilinear setting. For $1<p_1,p_2<\infty,\mathbf{P}=(p_1,p_2),0<\alpha<2n,\frac{\alpha}{n}<\frac{1}{p_1}+\frac{1}{p_2}$, and q such that $\frac{1}{q}=\frac{1}{p_1}+\frac{1}{p_2}-\frac{\alpha}{n}$, a vector weight $\omega=(\omega_1,\omega_2)$ belongs to $\mathbf{A_{P,q}}$ if

$$\sup_Q\left(\frac{1}{|Q|}\int_Q\mu_\omega(x)dx\right)\left(\frac{1}{|Q|}\int_Q\omega_1(x)^{-p_1'}dx\right)^{q/p_1'}\left(\frac{1}{|Q|}\int_Q\omega_2(x)^{-p_2'}dx\right)^{q/p_2'}<\infty.$$

where the notation $\mu_\omega=\omega_1^q\omega_2^q$. It was shown by Moen in [16] that if $\omega\in\mathbf{A_{P,q}}$ then $\omega_i^{-p_i'}\in A_{2p_i'}$ and $\mu_\omega\in A_{2q}$. In addition, the weights in $\mathbf{A_{P,q}}$ are precisely those for which

$$I_\alpha:L^{p_1}(\omega_1^{p_1})\times L^{p_2}(\omega_2^{p_2})\to L^q(\mu_\omega)$$

is bounded.

Now we return to our main results.

Theorem 1.1. *Let $1 < p, p_1, p_2, q < \infty, 0 < \alpha < n$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ and $\frac{1}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{\alpha}{n}$. If $[b, B_\alpha]_1$ is a compact operator from $L^{p_1} \times L^{p_2}$ to L^q, then $b \in$ CMO.*

Theorem 1.2. *Let $1 < p, p_1, p_2, q < \infty, 0 < \alpha < 2n$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, $\frac{\alpha}{n} < \frac{1}{p_1} + \frac{1}{p_2}$ and $\frac{1}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{\alpha}{n}$. For the local integral function b and $\vec{b} = (b, b)$, the following are equivalent,*

(A1) $b \in$ CMO.

(A2) $[\Pi\vec{b}, I_\alpha] : L^{p_1}(\omega_1^{p_1}) \times L^{p_2}(\omega_2^{p_2}) \to L^q(\omega_1^q \omega_2^q)$ *is a compact operator for all $\omega = (\omega_1, \omega_2)$ such that $\omega_1^{\frac{p_1 q}{p}}, \omega_2^{\frac{p_2 q}{p}} \in A_p$.*

(A3) $[\Pi\vec{b}, I_\alpha] : L^{p_1} \times L^{p_2} \to L^q$ *is a compact operator.*

2. Main lemmas

As mentioned in the introduction, CMO is the closure in BMO of the space of C^∞ functions with compact support. In [21], it was shown that CMO can be characterized in the following way.

Lemma 2.1. *([21]) Let $b \in$ BMO. Then b is in* CMO *if and only if*

$$\lim_{a \to 0} \sup_{|Q| = a} \frac{1}{|Q|} \int_Q |b(x) - b_Q| dx = 0; \tag{2.1}$$

$$\lim_{a \to \infty} \sup_{|Q| = a} \frac{1}{|Q|} \int_Q |b(x) - b_Q| dx = 0; \tag{2.2}$$

$$\lim_{|y| \to 0} \frac{1}{|Q|} \int_Q |b(x + y) - b_Q| dx = 0, \ \textit{for each } Q. \tag{2.3}$$

To prove Theorem 1.1 and Theorem 1.2, we need the following results.

Lemma 2.2. *Support that $b \in$ BMO with $\|b\|_* = 1$. If for some $0 < \epsilon < 1$ and a cube Q with its center at x_Q and r_Q, b is not a constant on cube Q and satisfies*

$$\frac{1}{|Q|} \int_Q |b(y) - b_Q| dy > \epsilon, \tag{2.4}$$

then for the functions $g(y) = \frac{|Q| \chi_{(2Q)^c}(y)}{|y - x_Q|^{n/p_2 + n}}$ and f is defined by

$$f(y) = |Q|^{-1/p_1} \big(\operatorname{sgn}(b(y) - b_Q)\big) \chi_Q(y), \tag{2.5}$$

There exists constants $\gamma_1, \gamma_2, \gamma_3$ satisfying $\gamma_2 > \gamma_1 > 2$ and $\gamma_3 > 0$, such that

$$\int_{\gamma_1 r_Q < |x-x_Q| < \gamma_2 r_Q} \big|[b, B_\alpha]_1(f,g)(x)\big|^q dx \geq \gamma_3^q, \tag{2.6}$$

$$\int_{|x-x_Q| > \gamma_2 r_Q} \big|[b, B_\alpha]_1(f,g)(x)\big|^q dx \leq \frac{\gamma_3^q}{4^q}. \tag{2.7}$$

Moreover, there exists a constant $0 < \beta << \gamma_2$ depending only on p_1, p_2, n such that for all measurable subsets $E \subset \big\{x : \gamma_1 r_Q < |x - x_Q| < \gamma_2 r_Q\big\}$ satisfying $\frac{|E|}{|Q|} < \beta^n$, we have

$$\int_E \big|[b, B_\alpha]_1(f,g)(x)\big|^q dx \leq \frac{\gamma_3^q}{4^q}. \tag{2.8}$$

Proof. It is easy to check that f satisfies

$$\begin{gathered}
\mathrm{supp} f \subset Q, \\
f(y)(b(y) - b_Q) = |Q|^{-1/p_1} |b(y) - b_Q| \chi_Q(y) \geq 0, \\
|f(y)| \leq |Q|^{-1/p_1}, \\
\|f\|_{L^{p_1}} \leq 1, \\
\int \big(b(y) - b_Q\big) f(y) dy = |Q|^{-1/p_1} \int_Q |b(y) - b_Q| dy,
\end{gathered}$$

and g satisfies that $\|g\|_{L^{p_2}} = C$ and for $x \in (2nQ)^c, y \in Q$, we get

$$|2x - y - y| = 2|x - y| \geq 2|x - x_Q| - 2|y - x_Q| \geq 2nr_Q - \sqrt{n} r_Q \geq n r_Q,$$

which implies that $2x - y \in \big(Q(y, 2\sqrt{n} r_Q)\big)^c \subset (2Q)^c$ and $g(2x - y) \approx |Q| \cdot |x - x_Q|^{-n/p_2 - n}$.

We first establish the following several technical estimates. For a cube Q with center x_Q and satisfying (2.4) for some $\epsilon > 0$ and $x \in (2nQ)^c$, the following point-wise estimates hold:

$$|B_\alpha\big((b - b_Q) f, g\big)(x)| \lesssim |Q|^{\frac{1}{p_1'} + 1} |x - x_Q|^{-2n - n/p_2 + \alpha}, \tag{2.9}$$

$$|B_\alpha\big((b - b_Q) f, g\big)(x)| \gtrsim \epsilon |Q|^{\frac{1}{p_1'} + 1} |x - x_Q|^{-2n - n/p_2 + \alpha}, \tag{2.10}$$

$$|B_\alpha\big(f, g\big)(x)| \lesssim |Q|^{\frac{1}{p_1'} + 1} |x - x_Q|^{-2n - n/p_2 + \alpha}, \tag{2.11}$$

where f, g as above and the constants involved are independent of b, f, g and ϵ.

To prove (2.9), from the fact that $\|b\|_* = 1$ and $x \in (2nQ)^c$, we have

$$|B_\alpha\big((b - b_Q) f, g\big)(x)| \;=\; \left| \int_Q \frac{(b(y) - b_Q) f(y) g(2x - y)}{|x - y|^{n - \alpha}} dy \right|$$

$$\begin{aligned} &\lesssim |Q|^{\frac{1}{p_1'}}|x-x_Q|^{-2n-n/p_2+\alpha}\int_Q (b(y)-b_Q)f(y)dy \\ &\lesssim |Q|^{\frac{1}{p_1'}+1}|x-x_Q|^{-2n-n/p_2+\alpha}. \end{aligned}$$

For (2.10), by $x \in (2nQ)^c$ and $y \in Q$, we have $|x-y| \approx |x-x_Q|$. Using that $\big(b(y)-b_Q\big)f(y) \geq 0$, we can compute

$$\begin{aligned} |B_\alpha\big((b-b_Q)f,g\big)(x)| &= \left|\int_Q \frac{(b(y)-b_Q)f(y)g(2x-y)}{|x-y|^{n-\alpha}}dy\right| \\ &\gtrsim |Q||x-x_Q|^{-2n-n/p_2+\alpha}\int_Q \big(b(y)-b_Q\big)f(y)dy \\ &= |Q|^{1/p_1'}|x-x_Q|^{-2n-n/p_2+\alpha}\int_Q \big|b(y)-b_Q\big|dy \\ &\gtrsim \epsilon|Q|^{\frac{1}{p_1'}+1}|x-x_Q|^{-2n-n/p_2+\alpha}. \end{aligned}$$

Finally using that $|f(y)| \leq |Q|^{-1/p_1}$ we obtain (2.20) as follows.

$$|B_\alpha\big(f,g\big)(x)| = \left|\int_Q \frac{f(y)g(2x-y)}{|x-y|^{n-\alpha}}dy\right| \lesssim |Q|^{\frac{1}{p_1'}+1}|x-x_Q|^{-2n-n/p_2+\alpha}.$$

Now, we give the proofs of (2.6)-(2.7). Note that for $b \in \mathrm{BMO}$, we have

$$\left(\int_{2^s r_Q<|x-x_Q|<2^{s+1}d_j} |b(x)-b_Q|^q dx\right)^{1/q} \lesssim s2^{sn/q}|Q|^{1/q}.$$

Taking $\nu > 16$, by (2.9) we obtain

$$\begin{aligned} &\left(\int_{|x-x_Q|>\nu r_Q} \big|(b(x)-b_Q)B_\alpha(f,g)(x)\big|^q dx\right)^{\frac{1}{q}} \\ &\leq C|Q|^{\frac{1}{p_1'}+1}\sum_{s=\lfloor \log_2 \nu\rfloor}^{\infty}\left(\int_{2^s r_Q<|x-x_Q|<2^{s+1}r_Q} \frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+n/p_2)}}dx\right)^{\frac{1}{q}} \\ &\leq C|Q|^{\frac{1}{p_1'}+1}\sum_{s=\lfloor \log_2 \nu\rfloor}^{\infty} 2^{-s(2n-\alpha+n/p_2)}|Q|^{-2+\frac{\alpha}{n}-\frac{1}{p_2}}\left(\int_{2^s r_Q<|x-x_Q|<2^{s+1}d_j} |b(x)-b_Q|^q dx\right)^{\frac{1}{q}} \\ &\leq C\sum_{s=\lfloor \log_2 \nu\rfloor}^{\infty} s2^{-s(2n-\alpha+\frac{n}{p_2}-\frac{n}{q})} \\ &\leq C\sum_{s=\lfloor \log_2 \nu\rfloor}^{\infty} 2^{-s(2n-\alpha+\frac{n}{p_2}-\frac{n}{q}-\frac{1}{2})} \end{aligned}$$

$$\leq C\nu^{-(2n-\alpha+\frac{n}{p_2}-\frac{n}{q}-\frac{1}{2})},$$

where we have used that $2n-\alpha+\frac{n}{p_2}-\frac{n}{q}-\frac{1}{2}>0$ and $s\leq 2^{s/2}$ for $4\leq\lfloor\log_2\nu\rfloor\leq s$.

For $\mu>\nu$, using (2.10) and the estimates above, we get

$$\begin{aligned}
&\int_{\nu r_Q<|x-x_Q|<\mu r_Q}\big|[b,B_\alpha](f,g)(x)\big|^q dx\\
&\geq C\bigg(\big|B_\alpha((b-b_Q)f,g)(x)\big|^q dx\bigg)^{1/q}\\
&\qquad -C\bigg(\int_{\nu r_Q<|x-x_Q|}\big|(b(x)-b_Q)B_\alpha(f,g)(x)\big|^q dx\bigg)^{1/q}\\
&\geq C\epsilon|Q|^{\frac{1}{p_1'}+1}\bigg(\int_{\nu r_Q<|x-x_Q|<\mu r_Q}|x-x_Q|^{q(-2n-n/p_2+\alpha)}dx\bigg)^{1/q}-C\nu^{-(2n-\alpha+\frac{n}{p_2}-\frac{n}{q}-\frac{1}{2})}\\
&\geq C\epsilon\Big(\nu^{-2nq+nq/p_2+\alpha q}-\mu^{-2nq+nq/p_2+\alpha q}\Big)^{\frac{1}{q}}-C\nu^{-(2n-\alpha+\frac{n}{p_2}-\frac{n}{q}-\frac{1}{2})}.
\end{aligned}$$

Once again, the constants appearing above are independent of Q. It is easy to see that we can select γ_1,γ_2 in place of ν,μ with $\gamma_2>>\gamma_1$, then (2.13) and (2.14) are verified for some $\gamma_3>0$.

We now verified (2.8). Let $E\subset\big\{\gamma_1 r_Q<|x-x_Q|<\gamma_2 r_Q\big\}$ be an arbitrary measurable set. It follows from Minkowski inequality that

$$\begin{aligned}
&\bigg(\int_E\big|[b,B_\alpha]_1(f,g)(x)\big|^q dx\bigg)^{\frac{1}{q}}\\
&\lesssim|Q|^{\frac{1}{p_1'}+1}\bigg(\int_E|x-x_Q|^{-q(2n+n/p_2-\alpha)}dx\bigg)^{\frac{1}{q}}+|Q|^{\frac{1}{p_1'}+1}\bigg(\int_E\frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+n/p_2)}}dx\bigg)^{\frac{1}{q}}\\
&\lesssim\bigg(\frac{|E|^{1/q}}{|Q|^{1/q}}+\Big(\frac{1}{|Q|}\int_E|b(x)-b_Q|^q dx\Big)^{\frac{1}{q}}\bigg)
\end{aligned}$$

It is proved in [6, p.269] that

$$\frac{1}{|Q|}\int_E|b(x)-b_Q|^q dx\lesssim\frac{|E|}{|Q|}\bigg(1+\log\Big(\frac{\tilde{C}|Q|}{|E|}\Big)\bigg)^{[q]+1}.$$

Taking $0<\beta<\min\{\tilde{C}^{1/n},\gamma_2\}$ and sufficiently small, then (2.8) holds. □

Lemma 2.3. *Support that $b\in$ BMO with $\|b\|_*=1$. If for some $0<\epsilon<1$ and a cube Q with its center at x_Q and r_Q, b is not a constant on cube Q and satisfies*

$$\frac{1}{|Q|}\int_Q|b(y)-b_Q|dy>\epsilon^{1/2},$$

then for the function $f_i(i=1,2)$ *defined by*

$$f_i(y_i)=|Q|^{-1/p_i}\big(sgn(b(y_i)-b_Q)-c_0\big)\chi_Q(y_i), \tag{2.12}$$

where $c_0=|Q|^{-1}\int_Q sgn\big(b(y)-b_Q\big)dy_i$ *and* $i=1,2$. *There exists constants* $\gamma_1,\gamma_2,\gamma_3$ *satisfying* $\gamma_2>\gamma_1>2$ *and* $\gamma_3>0$, *such that*

$$\int_{\gamma_1 r_Q<|x-x_Q|<\gamma_2 r_Q}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\geq\gamma_3^q, \tag{2.13}$$

$$\int_{|x-x_Q|>\gamma_2 r_Q}\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\leq\frac{\gamma_3^q}{4^q}. \tag{2.14}$$

Moreover, there exists a constant $0<\beta<<\gamma_2$ *depending only on* p_1,p_2,n *such that for all measurable subsets* $E\subset\big\{x:\gamma_1 r_Q<|x-x_Q|<\gamma_2 r_Q\big\}$ *satisfying* $\frac{|E|}{|Q|}<\beta^n$, *we have*

$$\int_E\big|[\Pi\vec{b},I_\alpha](f_1,f_2)(x)\big|^q dx\leq\frac{\gamma_3^q}{4^q}. \tag{2.15}$$

Proof. Since $\int_Q\big(b(y)-b_Q\big)dy=0$, it is easy to check that f_i satisfies

$$\begin{gathered}\mathrm{supp} f_i\subset Q,\\ f_i(y_i)(b(y)-b_Q)\geq 0,\\ \int f_i(y_i)dy_i=0,\\ |f_i(y_i)|\leq 2|Q|^{-1/p_i},\\ \|f_i\|_{L^{p_i}}\leq 2,\\ \int\big(b(y)-b_Q\big)f_i dy=|Q|^{-1/p_i}\int_Q|b(y_i)-b_Q|dy.\end{gathered}$$

For a cube Q with center x_Q and $x\in(2\sqrt{n}Q)^c$, the following point-wise estimates hold:

$$|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)|\lesssim|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}|x-x_Q|^{-2n+\alpha}, \tag{2.16}$$

$$|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)|\gtrsim\epsilon|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}|x-x_Q|^{-2n+\alpha}, \tag{2.17}$$

$$|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)|\lesssim|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{2.18}$$

$$|I_\alpha\big(f_1,(b-b_Q)f_2\big)(x)|\lesssim|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{2.19}$$

$$|I_\alpha\big(f_1,f_2\big)(x)|\lesssim|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}, \tag{2.20}$$

where f_i as above and the constants involved are independent of b, f_i and ϵ.

To prove (2.16), from the fact that $\|b\|_* = 1$ and $x \in (2\sqrt{n}Q)^c$, we have

$$\begin{aligned}
&|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)| \\
&= \left| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)(b(y_2)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \right| \\
&\lesssim |Q|^{-\frac{1}{p_1}-\frac{1}{p_2}}|x-x_Q|^{-2n+\alpha}\prod_{i=1}^2\int_Q (b(y_i)-b_Q)f_i(y_i)dy_i \\
&\lesssim |Q|^{-\frac{1}{p_1}-\frac{1}{p_2}}|x-x_Q|^{-2n+\alpha}\prod_{i=1}^2\int_Q |b(y_i)-b_Q|dy_i \\
&\lesssim |Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}|x-x_Q|^{-2n+\alpha}.
\end{aligned}$$

For (2.17), using that $\big(b(y_i)-b_Q\big)f_i(y_i) \geq 0$, we can compute

$$\begin{aligned}
&|I_\alpha\big((b-b_Q)f_1,(b-b_Q)f_2\big)(x)| \\
&= \left| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)(b(y_2)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \right| \\
&\gtrsim |x-x_Q|^{-2n+\alpha}\prod_{i=1}^2\left|\int_Q \big(b(y_i)-b_Q\big)f_i(y_i)dy_i\right| \\
&= |x-x_Q|^{-2n+\alpha}\prod_{i=1}^2 |Q|^{-\frac{1}{p_i}}\int_Q \big|b(y_i)-b_Q\big|dy_i \\
&\gtrsim \epsilon|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}|x-x_Q|^{-2n+\alpha}.
\end{aligned}$$

For (2.18), by the fact $|f_2(y_2)| \leq 2|Q|^{-1/p_2}$ and $\int_Q f_2(y_2)dy_2 = 0$, we can also estimate

$$\begin{aligned}
&|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)| \\
&= \left| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \right| \\
&= \left| \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}} dy_1dy_2 \right. \\
&\qquad \left. - \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(b(y_1)-b_Q)f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2'|^2\big)^{n-\alpha/2}} dy_1dy_2 \right| \\
&\lesssim |Q|^{-\frac{1}{p_1}-\frac{1}{p_2}}\int_Q\int_Q \frac{|y_2-y_2'|(b(y_1)-b_Q)f_1(y_1)}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}} dy_1dy_2
\end{aligned}$$

$$\lesssim |Q|^{-\frac{1}{p_1}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}\int_Q |b(y_1)-b_Q|dy_1$$
$$\lesssim |Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}.$$

It is easy to see that $|I_\alpha\big((b-b_Q)f_1,f_2\big)(x)|=|I_\alpha\big(f_1,(b-b_Q)f_2\big)(x)|$, then (2.19) holds. Finally using that f_1 has mean zero we obtain (2.20) as follows.

$$\begin{aligned}
&|I_\alpha\big(f_1,f_2\big)(x)|\\
&=\Big|\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{f_1(y_1)f_2(y_2)}{\big(|x-y_1|^2+|x-y_2|^2\big)^{n-\alpha/2}}-\frac{f_1(y_1)f_2(y_2)}{\big(|x-y_1'|^2+|x-y_2|^2\big)^{n-\alpha/2}}dy_1dy_2\Big|\\
&\lesssim |Q|^{-\frac{1}{p_1}-\frac{1}{p_2}}\int_Q\int_Q\frac{|y_1-y_1'||f_1(y_1)||f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2\\
&\lesssim |Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}|x-x_Q|^{-2n+\alpha-1}.
\end{aligned}$$

Now, we give the proofs of (2.13)-(2.15). Taking $\nu>16$, by (2.18) we obtain

$$\begin{aligned}
&\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)I_\alpha((b-b_Q)f_1,f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\\
&\le C|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}\Big(\int_{2^sr_Q<|x-x_Q|<2^{s+1}r_Q}\frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+1)}}dx\Big)^{\frac{1}{q}}\\
&\le C|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}2^{-s(2n-\alpha+1)}|Q|^{-2+\frac{\alpha-1}{n}}\Big(\int_{2^sr_Q<|x-x_Q|<2^{s+1}d_j}|b(x)-b_Q|^q dx\Big)^{\frac{1}{q}}\\
&\le C\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}s2^{-s(2n-\alpha+1-\frac{n}{q})}\\
&\le C\sum_{s=\lfloor\log_2\nu\rfloor}^{\infty}2^{-s(2n-\alpha-\frac{n}{q}+\frac{1}{2})}\\
&\le C\nu^{-(2n-\alpha-\frac{n}{q}+\frac{1}{2})},
\end{aligned}$$

where we have used that $s\le 2^{s/2}$ for $4\le\lfloor\log_2\nu\rfloor\le s$.

Similarly, we also have

$$\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)I_\alpha(f_1,(b-b_Q)f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\le C\nu^{-2n-\alpha+1-\frac{n}{q}},$$

$$\Big(\int_{|x-x_Q|>\nu r_Q}\big|(b(x)-b_Q)^2I_\alpha(f_1,f_2)(x)\big|^q dx\Big)^{\frac{1}{q}}\le C\nu^{-2n-\alpha+1-\frac{n}{q}}.$$

Then for $\mu > \nu$, using (2.16), (2.17) and the estimates above, we get

$$\begin{aligned}
&\int_{\nu r_Q<|x-x_Q|<\mu r_Q} \left|[\Pi\vec{b}, I_\alpha](f_1,f_2)(x)\right|^q dx\\
&\geq C\bigg(\left|I_\alpha((b-b_Q)f_1,(b-b_Q)f_2)(x)\right|^q dx\bigg)^{1/q}\\
&\quad -C\bigg(\int_{\nu r_Q<|x-x_Q|} \left|(b(x)-b_Q)I_\alpha((b-b_Q)f_1,f_2)(x)\right|^q dx\bigg)^{1/q}\\
&\quad -C\bigg(\int_{\nu r_Q<|x-x_Q|} \left|(b(x)-b_Q)I_\alpha(f_1,(b-b_Q)f_2)(x)\right|^q dx\bigg)^{1/q}\\
&\quad -C\bigg(\int_{\nu r_Q<|x-x_Q|} \left|(b(x)-b_Q)^2 I_\alpha(f_1,f_2)(x)\right|^q dx\bigg)^{1/q}\\
&\geq C\epsilon|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}\bigg(\int_{\nu r_Q<|x-x_Q|<\mu r_Q}|x-x_Q|^{q(-2n+\alpha)}dx\bigg)^{1/q} - C\nu^{-2n+\alpha+\frac{n}{q}-\frac{1}{2}}\\
&\geq C\epsilon\Big(\nu^{-2nq+n+\alpha q}-\mu^{-2nq+n+\alpha q}\Big)^{\frac{1}{q}} - C\nu^{-2n+\alpha+\frac{n}{q}-\frac{1}{2}}.
\end{aligned}$$

We can select γ_1, γ_2 in place of ν, μ with $\gamma_2 >> \gamma_1$, then (2.13) and (2.14) are verified for some $\gamma_3 > 0$.

We now verified (2.15). Let $E \subset \{\gamma_1 r_Q < |x-x_Q| < \gamma_2 r_Q\}$ be an arbitrary measurable set. It follows from Minkowski inequality that

$$\begin{aligned}
&\bigg(\int_E \left|[\Pi\vec{b}, I_\alpha](f_1,f_2)(x)\right|^q dx\bigg)^{\frac{1}{q}}\\
&\lesssim |Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}}\bigg(\int_E |x-x_Q|^{-q(2n-\alpha)}dx\bigg)^{\frac{1}{q}} + |Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}\bigg(\int_E \frac{|b(x)-b_Q|^q}{|x-x_Q|^{q(2n-\alpha+1)}}dx\bigg)^{\frac{1}{q}}\\
&\quad +|Q|^{\frac{1}{p_1'}+\frac{1}{p_2'}+\frac{1}{n}}\bigg(\int_E \frac{|b(x)-b_Q|^{2q}}{|x-x_Q|^{q(2n-\alpha+1)}}dx\bigg)^{\frac{1}{q}}\\
&\lesssim \bigg(\frac{|E|^{1/q}}{|Q|^{1/q}} + \Big(\frac{1}{|Q|}\int_E |b(x)-b_Q|^q dx\Big)^{\frac{1}{q}} + \Big(\frac{1}{|Q|}\int_E |b(x)-b_Q|^{2q}dx\Big)^{\frac{1}{q}}\bigg)
\end{aligned}$$

The same estimate as Lemma 2.2 and taking $0 < \beta < \min\{\tilde{C}^{1/n}, \gamma_2\}$, we can obtain the desired result. □

In the proof of the boundedness of the iterated commutators, the following two important properties of the weights we will be using.

Lemma 2.4. *([4]) Let $1 < p_1, p_2 < \infty, P = (p_1, p_2), 0 < \alpha < 2n, \frac{\alpha}{n} < \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$ and q such that $\frac{1}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{\alpha}{n}$. Suppose that $\omega_1^{\frac{p_1 q}{p}}, \omega_2^{\frac{p_2 q}{p}} \in A_p$. Then,*

(i) $\omega = (\omega_1, \omega_2) \in A_{P,q}$;

(ii) $\mu_\omega = \omega_1^q \omega_2^q \in A_p \subset A_q$.

In the proof of Theorem 1.2, we need the following weighted version of the Frechét-Kolmogorov-Riesz theorem. We refer to works by Hanche-Olsen and Holden [11] and Clop and Cruz [7].

Lemma 2.5. *Let $1 < q < \infty$ and $\omega \in A_q$. Suppose that the subset $\mathfrak{F} \subset L^q(\omega)$ satisfies the following conditions:*

(i) norm boundedness uniformly

$$\sup_{f \in \mathfrak{F}} \|f\|_{L^q(\omega)} < \infty; \tag{2.21}$$

(ii) translation continuity uniformly

$$\lim_{f \to 0} \|f(\cdot + y) - f(\cdot)\|_{L^q(\omega)} = 0 \ uniformly \ in \ f \in \mathfrak{F}; \tag{2.22}$$

(iii) control uniformly away from the origin

$$\lim_{A \to \infty} \int_{|x| > A} |f(x)|^q \omega(x) dx = 0 \ uniformly \ in \ f \in \mathfrak{F}; \tag{2.23}$$

then $\mathfrak{F}$ is pre-compact in $L^q(\omega)$.

Another reduction in the proof of Theorem 1.2 will be made by slightly modifying the bilinear fractional integral operator. This technique comes from Krantz and Li [14] (see also [1], [4]). More precisely, for any $\delta > 0$ small enough, the kernel $K^\delta(x, y_1, y_2)$ in $\mathbb{R}^{3n}$ such that for $\max\{|x - y_1|, |x - y_2|\} > 2\delta$,

$$K^\delta(x, y_1, y_2) = \frac{1}{\big(|x - y_1|^2 + |x - y_2|^2\big)^{n - \alpha/2}};$$

for $\max\{|x - y_1|, |x - y_2|\} < \delta$,

$$K^\delta(x, y_1, y_2) = 0;$$

and for all multi-indexes with $|\gamma| \leq 1$,

$$\partial^\gamma K^\delta(x, y_1, y_2) \lesssim \frac{1}{\big(|x - y_1| + |x - y_2|\big)^{2n - \alpha + |\gamma|}}.$$

The operators I_α^δ are defined by

$$I_\alpha^\delta(f_1,f_2)(x)=\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}K^\delta(x,y_1,y_2)f_1(y_1)f_2(y_2)dy_1dy_2.$$

Lemma 2.6. *If $b\in C_c^\infty$ and $\omega\in\mathbf{A_{P,q}}$, then*

$$\lim_\delta\big\|[\Pi\vec{b},I_\alpha^\delta]-[\Pi\vec{b},I_\alpha]\big\|_{L^{p_1}(\omega_1^{p_1})\times L^{p_1}(\omega_2^{p_2})\to L^q(\mu_\omega)}=0.$$

Proof. The proof of this result is very similar to that of Lemma 2.1 in [1] and we omit the details. □

3. Proofs of Theorem 1.1 and Theorem 1.2

Proof of Theorem 1.1. We first show that if $[b,B_\alpha]_1$ is bounded from $L^{p_1}\times L^{p_2}$ to L^p, then $b\in$ BMO. For $z_0\in\mathbb{R}^n\backslash\{0\}$, let $\delta=\frac{|z_0|}{2\sqrt{n}}$ and $Q_0(z_0,\delta)$ denote the open cube centered at z_0 with side length 2δ. Then $|x|^{n-\alpha}$ has an absolutely convergent Fourier series

$$|x|^{n-\alpha}=\sum a_me^{iv_m\cdot x}$$

with $\sum|a_m|<\infty$, where the exact form of the vectors v_m is unrelated. Taking $z_1=\frac{z_0}{\delta}$ we have the expansion

$$|x|^{n-\alpha}=\delta^{-n+\alpha}|\delta x|^{n-\alpha}=\delta^{-n+\alpha}\sum a_me^{iv_m\cdot\delta x}\ for\ |x-z_1|<\sqrt{n}.$$

Given cubes $Q=Q(x_0,r)$ and $Q'=Q(x_0-rz_1,r)$, if $x\in Q$ and $y\in Q'$, then

$$\Big|\frac{x-y}{r}-\frac{z_0}{\delta}\Big|\le\Big|\frac{x-x_0}{r}\Big|+\Big|\frac{y-(x_0-\frac{rz_0}{\delta})}{r}\Big|<\sqrt{n},$$

and

$$|2x-y-(x_0-rz_1)|\le|x-x_0|+|x-y-rz_1|\le\frac{\sqrt{n}}{2}r+r\Big|\frac{x-y}{r}-\frac{z_0}{\delta}\Big|\le\frac{3\sqrt{n}}{2}r,$$

which implies that $2x-y\in\tilde{Q}=3\sqrt{n}Q'$.

Let $s(x)=\overline{\operatorname{sgn}(\int_{Q'}(b(x)-b(y))dy)}$. Then

$$\begin{aligned}
&\frac{1}{|Q|}\int_Q|b(x)-b_{Q'}|dx\\
&=\frac{1}{|Q|}\frac{1}{|Q'|}\int_Q\Big|\int_{Q'}(b(x)-b(y))dy\Big|dx\\
&=\frac{1}{|Q|^2}\int_Q\int_{Q'}s(x)(b(x)-b(y))dydx\\
&=\frac{1}{|Q|^2}\int_Q\int_{Q'}\frac{r^{n-\alpha}s(x)(b(x)-b(y))}{|x-y|^{n-\alpha}}\Big|\frac{x-y}{r}\Big|^{n-\alpha}dydx
\end{aligned}$$

$$= \frac{1}{|Q|}\int_Q\int_{Q'} \frac{s(x)(b(x)-b(y))}{|x-y|^{n-\alpha}} \sum a_m e^{iv_m\cdot\delta\frac{x-y}{r}}dydx$$
$$= \frac{1}{|Q|}\int_Q\int_{Q'} \frac{s(x)(b(x)-b(y))}{|x-y|^{n-\alpha}} \sum a_m e^{iv_m\cdot\delta\frac{2x-y}{r}}e^{-iv_m\cdot\delta\frac{y}{r}}dydx$$
$$= \frac{1}{|Q|}\int_Q\int_{Q'} \frac{s(x)(b(x)-b(y))}{|x-y|^{n-\alpha}} \sum a_m e^{iv_m\cdot\delta\frac{2x-y}{r}}e^{-iv_m\cdot\delta\frac{y}{r}}\chi_Q(x)\chi_{Q'}(y)\chi_{\tilde{Q}}(2x-y)dydx$$

Setting

$$f_m(y) = e^{-iv_m\cdot\frac{\delta}{r}y}\chi_{Q'}(y),$$
$$g_m(z) = e^{-iv_m\cdot\frac{\delta}{r}z}\chi_{\tilde{Q}}(z),$$
$$h_m(x) = e^{iv_m\cdot\frac{\delta}{r}x}s(x)\chi_Q(x),$$

we have

$$\begin{aligned}\frac{1}{|Q|}\int_Q |b(x)-b_{Q'}|dx &= \frac{1}{|Q|}\sum a_m\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{(b(x)-b(y))f_m(y)g_m(2x-y)}{|x-y|^{n-\alpha}}h_m(x)dydx\\ &= \frac{1}{|Q|}\sum a_m\int_{\mathbb{R}^n}[b,B_\alpha]_1(f_m,g_m)(x)h_m(x)dx.\end{aligned}$$

It follows that

$$\begin{aligned}\frac{1}{|Q|}\int_Q |b(x)-b_Q|dx &\lesssim \frac{1}{|Q|}\int_Q |b(x)-b_{Q'}|dx\\ &\lesssim \frac{1}{|Q|^{1/q}}\sum |a_m|\|[b,B_\alpha]_1(f_m,g_m)\|_{L^q}\\ &\lesssim \frac{\|f_m\|_{L^{p_1}}\|g_m\|_{L^p_2}}{|Q|^{1/p}}\|[b,B_\alpha]_1\|_{L^{p_1}\times L^{p_2}\to L^q}\\ &\lesssim \|[b,B_\alpha]_1\|_{L^{p_1}\times L^{p_2}\to L^q},\end{aligned}$$

which yields $b\in$ BMO and $\|b\|_*\lesssim C\|[b,B_\alpha]_1\|_{L^{p_1}\times L^{p_2}\to L^q},$.

To prove b be an element of CMO, we will adapt some arguments from [5], see also [4], which in turn are based on the original work in [21]. The approach is the following: if one of the conditions Eqs.(2.1)-(2.3) in Lemma 2.1 is failed, we will show that there exist sequences of functions, $\{f_j\}_j$ uniformly bounded on L^{p_1} and $\{g_j\}_j$ uniformly bounded on L^{p_2}, such that $[b,B_\alpha]_1(f_j,g_j)$ has no convergent subsequence, which contradicts the assumption that $[b,B_\alpha]_1$ is compact. It gives us that if $[b,B_\alpha]_1$ is compact, b must satisfy all three conditions; that is $b\in$ CMO.

By Lemma 2.2, it is sufficient to once again repeat the steps preformed in [5] (or [4], [6]) to obtain the desired result. Hence, the proof of Theorem 1.1 is completed. □

Proof of Theorem 1.2. $(A1) \Rightarrow (A2)$: Note that (2.21) is immediate since for $b \in C_c^\infty$, $[b, I_\alpha^\delta]$ is bounded from $L^{p_1}(\omega_1^{p_1}) \times L^{p_2}(\omega_2^{p_2})$ to $L^q(\mu_\omega)$, because $\omega \in A_{P,q}$ by Lemma 2.6.

To show that (2.22) holds, we can use a similar method as in [4, p.491], the proof of this results is very similar to that of linear commutator case, we omit the detail.

Now we give the estimate for (2.23). Denote

$$\mathfrak{F} = \big\{[\Pi\vec{b}, I_\alpha^\delta](f_1, f_2) : f_i \in L^{p_i}(\omega_i^{p_i}), \|f_i\|_{L^{p_i}(\omega_i)} \le 1, i = 1, 2\big\}.$$

Then $\mathfrak{F}$ is uniformly bounded, because $[\Pi\vec{b}, I_\alpha^\delta]$ is a bounded operator from $L^{p_1}(\omega_1^{p_1}) \times L^{p_2}(\omega_2^{p_2})$ to $L^q(\mu_\omega)$, because $\omega \in A_{P,q}$ by Lemma 2.1. To prove the uniform equicontinuity of $\mathfrak{F}$, we must see that

$$\lim_{t\to 0} \|[\Pi\vec{b}, I_\alpha^\delta](f_1, f_2)(\cdot + t) - [\Pi\vec{b}, I_\alpha^\delta](f_1, f_2)(\cdot)\|_{L^q(\mu_\omega)} = 0.$$

To do this, we write

$$\begin{aligned}
&[\Pi\vec{b}, I_\alpha^\delta](f_1, f_2)(x) - [\Pi\vec{b}, I_\alpha^\delta](f_1, f_2)(x + t)\\
&= \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \big(b(x) - b(y_1)\big)\big(b(x) - b(y_2)\big)K^\delta(x, y_1, y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad - \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \big(b(x + t) - b(y_1)\big)\big(b(x + t) - b(y_2)\big)K^\delta(x + t, y_1, y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&= \big(b(x) - b(x + t)\big)^2 \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} K^\delta(x, y_1, y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad + \big(b(x) - b(x + t)\big) \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \big(b(x + t) - b(y_2)\big)K^\delta(x, y_1, y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad + \big(b(x) - b(x + t)\big) \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \big(b(x + t) - b(y_1)\big)K^\delta(x, y_1, y_2)f_1(y_1)f_2(y_2)dy_1dy_2\\
&\quad + \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \big(b(x + t) - b(y_1)\big)\big(b(x + t) - b(y_2)\big)\\
&\qquad\qquad \times \big(K^\delta(x, y_1, y_2) - K^\delta(x + t, y_1, y_2)\big)f_1(y_1)f_2(y_2)dy_1dy_2\\
&= I_1(x, t) + I_2(x, t) + I_3(x, t) + I_4(x, t).
\end{aligned}$$

For $I_1(x, t)$, we simply have

$$|I_1(x, t)| \lesssim |t|^2\|\nabla b\|_\infty I_\alpha(|f_1|, |f_2|)(x),$$

which implies that

$$\|I_1(\cdot, t)\|_{L^q(\omega)} \lesssim |t|^2.$$

Similarly, we also have that for $j = 2, 3$

$$|I_j(x, t)| \lesssim |t|\|\nabla b\|_\infty\|b\|_\infty I_\alpha(|f_1|, |f_2|)(x),$$

and

$$\|I_j(\cdot,t)\|_{L^q(\omega)} \lesssim |t|.$$

We now give the estimate for $I_4(x,t)$. We may assume that $|t| \in (0,\delta/4)$. Thus, if $\max\{|x-y_1|,|x-y_2|\} \le \delta/2$ we have

$$K^\delta(x+t,y_1,y_2) - K^\delta(x,y_1,y_2) = 0$$

and $\max\{|x-y_1|,|x-y_2|\} > \delta/2$ we have

$$\max\{|x-y_1|,|x-y_2|\} > 2t.$$

This gives us that

$$\begin{aligned}
\big|I_4(x,t)\big| &\lesssim |t|\|b\|_\infty^2 \int\int_{\max\{|x-y_1|,|x-y_2|\}>\delta/2} \frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2 \\
&\lesssim |t|\|b\|_\infty^2 \sum_{j\ge 0}\int\int_{2^{j-1}\delta<\max\{|x-y_1|,|x-y_2|\}<2^j\delta} \frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha+1}}dy_1dy_2 \\
&\lesssim |t|\|b\|_\infty^2 \sum_{j\ge 0}\int\int_{2^{j-1}\delta<\max\{|x-y_1|,|x-y_2|\}<2^j\delta} \frac{|f_1(y_1)f_2(y_2)|}{\big(|x-y_1|+|x-y_2|\big)^{2n-\alpha}} \\
&\qquad \times \frac{1}{\max\{|x-y_1|,|x-y_2|\}}dy_1dy_2 \\
&\lesssim \frac{|t|}{\delta}\|b\|_\infty^2 I_\alpha(|f_1|,|f_2|)(x),
\end{aligned}$$

which gives

$$\|I_4(\cdot,t)\|_{L^q(\mu_\omega)} \le |t|.$$

Combining the estimates above and let $|t|<1$, we have

$$\|[\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(\cdot+t) - [\Pi\vec{b},I_\alpha^\delta](f_1,f_2)(\cdot)\|_{L^q(\mu_\omega)} \lesssim |t|.$$

Thus, if $b\in$ CMO, $[\Pi\vec{b},I_\alpha]$ is a compact operator from $L^{p_1}(\omega_1^{p_1})\times L^{p_2}(\omega_2^{p_2})$ to $L^q(\mu_\omega)$.

Obviously $(A2)\Rightarrow(A3)$. So it remains to show that $(A3)\Rightarrow(A1)$. By Lemma 2.3 and the same argument as Theorem 1.1, we need only to prove $b\in$ BMO.

$(A3)\Rightarrow(A1)$: Let $z_0\in\mathbb{R}^n$ such that $|(z_0,z_0)|>2\sqrt{n}$ and let δ small enough such that $\delta<1$. Take $B=B\big((z_0,z_0),\delta\sqrt{2n}\big)\subset\mathbb{R}^{2n}$ be the ball for which we can express $(|y_1|^2+|y_2|^2)^{n-\alpha/2}$ as an absolutely convergent Fourier series of the form

$$(|y_1|^2+|y_2|^2)^{n-\alpha/2} = \sum_j a_j e^{iv_j\cdot(y_1,y_2)}, \quad (y_1,y_2)\in B,$$

where $\sum_j|a_j|<\infty$ and we do not care about the vectors $v_j\in\mathbb{R}^{2n}$, but we will at times express them as $v_j=(v_j^1,v_j^2)\in\mathbb{R}^n\times\mathbb{R}^n$.

Set $z_1 = \delta^{-1} z_0$ and note that

$$\big(|y_1 - z_1|^2 + |y_2 - z_1|^2\big)^{1/2} < \sqrt{2n} \Rightarrow \big(|\delta y_1 - z_0|^2 + |\delta y_2 - z_0|^2\big)^{1/2} < \delta\sqrt{2n}.$$

Then for any (y_1, y_2) satisfying the inequality on the left, we have

$$(|y_1|^2 + |y_2|^2)^{n-\alpha/2} = \delta^{-2n+\alpha}(|\delta y_1|^2 + |\delta y_2|^2)^{n-\alpha/2} = \delta^{-2n+\alpha} \sum_j a_j e^{i\delta v_j \cdot (y_1, y_2)}.$$

Let $Q = Q(x_0, r)$ be any arbitrary cube in $\mathbb{R}^n$. Set $\tilde{z} = x_0 + r z_1$ and take $Q' = Q(\tilde{z}, r) \subset \mathbb{R}^n$. So for any $x \in Q$ and $y_1, y_2 \in Q'$, we have

$$\Big|\frac{x - y_1}{r} - z_1\Big| \le \Big|\frac{x - x_0}{r}\Big| + \Big|\frac{y_1 - \tilde{z}}{r}\Big| \le \sqrt{n}, \quad \Big|\frac{x - y_2}{r} - z_1\Big| \le \Big|\frac{x - x_0}{r}\Big| + \Big|\frac{y_2 - \tilde{z}}{r}\Big| \le \sqrt{n},$$

which implies that

$$\left(\Big|\frac{x - y_1}{r} - z_1\Big|^2 + \Big|\frac{x - y_2}{r} - z_1\Big|^2\right)^{1/2} \le \sqrt{2n}.$$

Let $s(x) = \overline{\operatorname{sgn}(\int_{Q'}(b(x) - b(y))dy)}$. We have the following estimate,

$$\begin{aligned}
&\Big(\frac{1}{|Q|}\int_Q |b(x) - b_Q|dx\Big)^2 \lesssim \Big(\frac{1}{|Q|}\int_Q |b(x) - b_{Q'}|dx\Big)^2 \\
&\lesssim \frac{1}{|Q|}\int_Q |b(x) - b_{Q'}|^2 dx \lesssim \frac{1}{|Q|}\int_Q s(x)^2 (b(x) - b_{Q'})^2 dx \\
&\lesssim \frac{1}{|Q||Q'|^2}\int_Q\int_{Q'}\int_{Q'} s(x)^2 \big(b(x) - b(y_1)\big)\big(b(x) - b(y_2)\big) dy_1 dy_2 dx \\
&\lesssim \frac{r^{2n-\alpha}\delta^{-2n+\alpha}}{|Q|^3}\int_Q\int_{Q'}\int_{Q'} \frac{s(x)^2\big(b(x) - b(y_1)\big)\big(b(x) - b(y_2)\big)}{\big(|x - y_1|^2 + |x - y_2|^2\big)^{n-\alpha/2}} \sum_j a_j e^{i\frac{\delta}{r} v_j \cdot (x - y_1, x - y_2)} dy_1 dy_2 dx.
\end{aligned}$$

Setting

$$\begin{aligned}
g_j(y_1) &= e^{-i\frac{\delta}{r} v_j^1 \cdot y_1} \chi_{Q'}(y_1), \\
h_j(y_2) &= e^{-i\frac{\delta}{r} v_j^2 \cdot y_2} \chi_{Q'}(y_2), \\
m_j(x) &= e^{i\frac{\delta}{r} v_j \cdot (x, x)} \chi_Q(x) s(x)^2.
\end{aligned}$$

We have

$$\begin{aligned}
&\Big(\frac{1}{|Q|}\int_Q |b(x) - b_Q|dx\Big)^2 \\
&\lesssim \frac{r^{2n-\alpha}\delta^{-2n+\alpha}}{|Q|^3} \sum_j |a_j| \int_{\mathbb{R}^n} \big|[\Pi\vec{b}, I_\alpha](g_j, h_j)(x) m_j(x)\big| dx \\
&\lesssim r^{-n-\alpha}\delta^{-2n+\alpha} \sum_j |a_j| \|[\Pi\vec{b}, I_\alpha]\|_{L^{p_1} \times L^{p_2} \to L^q} \|g_j\|_{L^{p_1}} \|h_j\|_{L^{p_2}} \|m_j\|_{L^{q'}}
\end{aligned}$$

$$\lesssim \delta^{-2n+\alpha}\|[\Pi\vec{b}, I_\alpha]\|_{L^{p_1}\times L^{p_2}\to L^q}\sum_j |a_j|.$$

The desired result follows from here. □

Acknowledgments We would like to thank the anonymous referee for his/her comments.

References

[1] Bényi, Á., Damián, W., Moen, K., Torres, R.H.: Compactness properties of commutators of bilinear frctional integrals. *Math. Z.*, (2015), doi: 10.1007/s00209-015-1437-4.

[2] Bényi, Á., Damián, W., Moen, K., Torres, R.H.: Compact bilinear operators: the weighted case. Mich. Math. J. **4**, 39-51 (2015)

[3] Bényi, Á., Torres, R.H.: Compact bilinear operators and commutators. Proc. Amer. Math. Soc. **141**(10), 3609-3621 (2013)

[4] Chaffee, L., Torres, R.H.: Characterization of Compactness of the Commutators of Bilinear Fractional Integral Operators. Potential Anal. **43**, 481-494 (2015)

[5] Chen, Y., Ding, Y., Wang, X.: Compactness of commutators of Riesz potential on Morrey spaces. Potential Anal. **30**, 301-313(2009)

[6] Chen, Y., Ding, Y., Wang, X.: Compactness of commutators for Singular integrals on Morrey spaces. Canad. J. Math. **64**, 257-281(2012)

[7] Clop, A., Cruz, V.: Weighted estimates for Beltrami equations. Ann. Acad. Sci. Fenn. Math. **38**, 91-113 (2013)

[8] Coifman, R., Rochberg, R., Weiss,G.: Factorization theorems for Hardy spaces in several variables. *Ann. of Math.*, **103**, 611-635 (1976)

[9] Grafakos, L.: On multilinear fractional integrals. Studia Math. **102**, 49-56 (1992)

[10] L. Grafakos, L., Kalton, N.: Some remarks on multilinear maps and interpolation, Math. Ann. **319**, 151-180 (2001)

[11] Hanche-Olsen, H., Holden, H.: The Kolmogorov-Riesz compactness theorem. Expo. Math. **28**, 385-394 (2010)

[12] Kenig, C., Stein, E.: Multilinear estimates and fractional integration, Math. Res. Lett. **6** 1-15 (1999)

[13] Krantz, S., Li, S.-Y.: Boundedness and compactness of integral operators on spaces of homogeneous type and applications, I. J. Math. Anal. Appl. **258**, 629-641 (2001)

[14] Krantz, S., Li, S.-Y.: Boundedness and compactness of integral operators on spaces of homogeneous type and applications, II. J. Math. Anal. Appl. **258**, 642-657 (2001)

[15] Lerner, A.K., Ombrosi, S., Pérez, C., Torres, R.H., Trujillo-González, R.: New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory. *Adv. Math.*, **220**, 1222-1264 (2009)

[16] Moen, K.: Weighted inequalities for multilinear fractional integral operators. Collect Math. **60**, 213-238 (2009)

[17] Pérez, C.: Endpoint estimates for commutators of singular integral operators. *J. Funct. Anal.*, **128**, 163-185 (1995)

[18] Pérez, C., Torres, R.H.: Sharp maximal function estimates for multilinear singular integrals. *Contemp. Math.*, **320**, 323-331 (2003)

[19] Stein, E.M.: Singular Integral and Differentiability Properties of Functions. Princeton University Press, Princeton (1971)

[20] Tang, L.: Weighted estimates for vector-valued commutators of multilinear operators. *Proc. Roy. Soc. Edinburgh Sect. A*, **138**, 897-922 (2008)

[21] Uchiyama, A.: On the compactness of operators of Hankel type. Tôhoku Math. J. **30**, 163-171(1978)

[22] Wang, S.: The compactness of the commutator of fractional integral operator (in Chinese). Chin. Ann. Math **8**(A), 475-482(1987)

College of Mathematics and System Sciences
Xinjiang University
Urumqi 830046
Republic of China
E-mail address: Wangdh1990@126.com; zhoujiangshuxue@126.com

School of Mathematics and Statistics
WuHan University
WuHan 430072
Republic of China
E-mail address: wychencn@hotmail.com